

Aggressive Exercise as Treatment for Chronic Low Back Pain

Isaac Cohen¹ and James Rainville^{2,3}

- 1 The Orthopaedic and Sports Medicine Center, Trumbull, Connecticut, USA
- 2 The Spine Center, New England Baptist, Boston, Massachusetts, USA
- 3 Department of Physical Medicine and Rehabilitation, Harvard Medical School, Boston, Massachusetts, USA

Contents

Abstract	75
1. Quantification of Back Function	76
1.1 Flexibility	76
1.2 Trunk Strength	77
1.3 Lifting Capacity	77
1.4 Cardiovascular Endurance	77
2. Patient Education and Behavioural Management	77
3. Exercise Regimens	78
4. Studies Utilising Aggressive Exercise Regimens	79
5. Conclusion	79

Abstract

Exercise has long been a standard of treatment for back pain. Over the last 2 decades, the use of intense, non-pain-contingent exercises for treatment of chronic back pain has received increased advocacy. The main goals of these treatments are to improve functioning of painful lumbar soft tissue and to decrease the fears and concerns of patients about using their backs for daily activities. The methodology of an aggressive quota-based exercise approach to back pain is outlined in this article. This approach relies on objective quantification of physical capabilities, treatment directed at altering these parameters, and repeat quantification for determination of treatment efficacy and positive feedback. By eliminating impairments in back function, altering fears and beliefs about pain, and reducing disability, patients with chronic low back pain can achieve meaningful improvements in their quality of life.

Chronic back pain is a complicated medical predicament. Pain is a subjective phenomenon. The perception and subsequent reporting of pain is influenced by cognitive and social factors in addition

to the peripheral pain stimulus. These factors may influence self-report of both pain and disability, and influence responsiveness to treatments.^[1-4] Many individuals with chronic back pain have strong pain

beliefs,^[5,6] low self-efficacy, and are fearful that activities may increase pain or cause further injury.^[7,8] These cognitive factors strongly influence the functional level of the individual with back pain. Additionally, many healthcare providers advise patients to avoid activities that stimulate back pain, and thereby reinforce activity restrictions.^[9] Loss of physical capacities, or deconditioning, occurs as a consequence of persistent activity restrictions. These physical impairments result in disability, or decreased ability to engage in activities of daily living.

The multidimensional nature of chronic low back pain is best addressed with a comprehensive approach that recognises and treats deconditioning and detrimental cognitive behavioural elements. Pain-focused treatments neglect these factors and are insufficient to alter patients' complex predicaments.

As a therapeutic modality, exercise has a primary goal of improving functioning of targeted tissues, that is, tissue length, tissue resilience, muscle strength and endurance. From a cognitive standpoint, successful completion of exercise in the presence of chronic pain lessens patients' fears and concerns, improves self-efficacy and confidence for performing daily activities, thereby decreasing disability.^[5,10]

Many healthcare providers are cautious about recommending exercises or any activities that may elicit back pain for fear that increased pain would indicate harm to degenerative spinal structures.^[9,11] This reluctance is perhaps the most important factor limiting the prescription of aggressive exercise for treatment of chronic back pain. Although a growing body of literature supports the safety and efficacy of aggressive exercise,^[12-21] providers' scepticism must be overcome to utilise this approach. It is important that all providers involved with the care of these patients, hold beliefs that normal back function can occur in the presence of chronic back pain, and that symptoms produced with exercise are generally benign. Without these common be-

liefs, an aggressive exercise approach is unlikely to succeed.

This approach is indicated for conditions such as lumbar disc degeneration, herniated nucleus pulposus, spinal stenosis, facet syndrome, Grade I-II spondylolisthesis, spondylolysis, myofascial pain and postoperative status (fusion, laminectomy). This approach is contraindicated in medical instability, lesions best treated with surgery, severe osteoporosis, fracture, tumour, cauda equina and conus medullaris syndromes, progressive neurological deficit, spinal instability, Grade III-IV spondylolisthesis, visceral/systemic pathology and spondyloarthropathy.

1. Quantification of Back Function

The goal of aggressive exercise is to improve back function. The cornerstone of this approach requires that back function be quantified so that progress towards this goal can be measured. In general, this includes measurements of trunk and lower extremity flexibility, trunk strength, lifting capacity and cardiovascular endurance. It is important to be aware that a patient's performance may be limited by pain, illness behaviours, and fears of re-injury or exacerbating their symptoms. In this sense, performance is a psychophysical phenomenon and not a purely physical event. Quantification initially serves to identify baseline impairments and guide clinicians in the selection of an appropriate level of exercise.

1.1 Flexibility

Deficits in flexibility in patients with chronic back pain have been demonstrated in multiple studies.^[22-27] The motions assessed include total lumbosacral flexion, extension, side bending and straight leg raising. Total lumbosacral sagittal motion assesses both lumbar and pelvic components, and straight leg raising reflects hamstring flexibility. Multiple techniques have been described in the literature to assess lumbar spine motion, such as the Schober (tape measure) test and its modifications, flexicurves, goniometers, finger-to-floor test, in-

clinometers, spinoscopy and radiographic methods.^[24,27-33] In the authors' practice, the single inclinometer technique is used, as it is simple, reliable and easy to interpret.^[34] Normal values for trunk motion have been established: total flexion 100 to 120°, extension 25 to 45°, side bending 25 to 45°, and straight leg raising 75 to 85°.^[23,24,26,35]

1.2 Trunk Strength

Studies have shown that patients with chronic low back pain have deficits in trunk strength.^[36-44] The loss of extensor strength is much greater than that of flexor strength.^[36,38,40,41] The normal extensor to flexor strength ratio is 1.2 to 1.5, and in patients with chronic back pain it has been documented to be less than 1.0.^[36,41] Although there are several methods available for documenting trunk strength (isometric, isokinetic and isoinertial), we prefer the isoinertial method, as it is more functional. This type of resistance is provided by free weights and most machines, and applies a constant load throughout the range of motion. At our institutions, Cybex Back Extension Machines (Lumex Corp, Ronkonkoma, NY) provide quantitative information regarding isoinertial capacity and allow for easy transference of exercise to a health club. Baseline assessment entails measuring the maximum amount of weight that a patient can correctly lift for four repetitions. The ideal is for patients to perform repetitions at 100% of ideal body-weight (IBW).^[45]

1.3 Lifting Capacity

Lifting capacity is included in the assessment of back function as it is a common activity of daily living and involves integration of lumbar function with other functional units. Deficits of isoinertial lifting capacity of 40 to 60% of normal have been associated with chronic low back pain.^[46,47] We utilise a standardised isoinertial lifting protocol, in which a patient lifts a plastic crate from floor to waist and waist to shoulder level with increasingly heavier loads.^[48] Patients' goals are to achieve normal lifting capacity: 50% IBW from floor to

waist (lumbar lift) and 40% IBW from waist to shoulder (cervical lift) for men, and 35% IBW floor to waist and 25% waist to shoulder for women.^[48]

1.4 Cardiovascular Endurance

The consequences of deconditioning on the cardiovascular system are well recognised.^[49-53] Less is known of the effects of deconditioning in patients with chronic back pain, but its existence has been documented.^[54,55] Cardiovascular conditioning can improve patients' tolerance for physical activities and may have beneficial effects on mood, sleep and muscle relaxation.^[55-59] We assess cardiovascular fitness with lower extremity ergometers (Cybex Fitron: Cybex Fitron, Ronkonkoma, NY, USA), which provide quantitative information in units of kg • m/min. The target heart rate is 75 to 85% of age-determined heart rate, or as determined by the referring physician for patients with cardiac disease. Testing is done using a 9-minute protocol. Patients are started on predetermined workloads which are sequentially increased every 3 minutes, using heart rate and psychophysical tolerance as endpoints. Although norms are not available for this equipment, total work of less than 7000 kg • m/min for men and 6000 kg • m/min for women on lower extremity testing is considered suboptimal.^[45]

2. Patient Education and Behavioural Management

Illness behaviours and counterproductive attitudes and beliefs are components of chronic low back pain.^[5-8,60-62] It is critical that they be addressed to improve the patient's predicament. Illness behaviours may be inadvertently rewarded with attention and concern from family and friends, health-care providers, or financially compensated with disability payments. During treatment, clinicians educate patients regarding their spine anatomy, pathology and physiology, and the deleterious effects of deconditioning. They are informed about the beneficial effects of exercise on their condition and

made aware that an initial increase in their pain is expected. They are also informed that exercise is generally safe, and that they are expected to continue to exercise in spite of an increase in symptoms. This information provides a rationale for exercise, establishes realistic treatment expectations and focuses the patients' efforts on impairments and disability as opposed to pain reduction.

Exercise recommendations are based on past performance levels and the patient's goals, and are not contingent on daily pain complaints. This 'quota-based' exercise programme is very successful for inducing physiological changes in back function. As exercise is based on measured performance, patients are continuously given numerical feedback regarding their increasing physical capacities. This provides tangible evidence that challenges their beliefs regarding pain and function.^[10]

Dramatic expressions of pain are acknowledged and evaluated, but the responses to these expressions are limited. Patients are advised to perform at their best possible level during these exacerbations, and attempts are made to keep the patient focused on treatment goals. Resumption of meaningful life activities is continuously encouraged by members of the team.

Although not essential, we have found that treating some patients in a group setting can be advantageous. Group sessions allow modelling, where patients experiencing success support those new or struggling during the rehabilitation process.

During treatment, patients are strongly encouraged to discontinue opioid agonists (narcotics), muscle relaxants and sedatives. Patients with significant depression or anxiety are given additional support and may be referred to psychologists or psychiatrists.

Weekly meetings of the treatment team are held to discuss patients' current level of function, treatment goals, medical and behavioural issues. It is crucial for the team of providers to be unified and provide a consistent message that normal back function may be safely re-established in spite of pain.

3. Exercise Regimens

Provided that there are no contraindications to aggressive exercise, patients are assigned to an exercise level based on measured impairments. During the initial therapy evaluation, instructions are provided regarding a home stretching programme specific for measured impairments in flexibility. Stretching at the physiological limits of flexibility is performed at least twice per day. Following assessment of strength, treatment goals are established according to a patient's age, gender and IBW. Patients are carefully instructed on the correct set-up and use of all exercise equipment. The next 1 to 2 weeks is dedicated towards acquiring proper exercise technique and diminishing fear of physical exertion. A list of exercises and targeted back muscles is provided in table A1 and figure A1.

The second phase of treatment generally lasts between 2 and 4 weeks, during which rapid progression towards treatment goals is expected. Patients generally meet with their therapist 2 to 3 times per week with sessions lasting between 1 and 2.5 hours. Patients who are highly motivated and experienced with exercise require only 1 weekly therapy session and exercise independently several more times per week. Exercise sessions involve supervised stretching, aerobic conditioning, general strengthening, specific back strengthening, and lifting. Repeat quantification is performed every 2 weeks to monitor progress, provide feedback and document treatment outcomes. Patients are encouraged to achieve their pre-established goals by increasing their repetitions and/or weights with each session. Exercise must provide sufficient physiologic overload to produce improvements in physical abilities. In the presence of chronic pain, this intensity of exercise will often stimulate abnormally sensitive nociceptors associated with the chronic pain symptoms and exacerbate pain complaints. These exacerbations are usually tolerable, brief and do not represent tissue damage. As training progresses, tissue function improves and the sensitivity of abnormal nociceptors tends to decrease. Therapy continues in spite of ongoing

pain complaints, and wellness behaviours are reinforced.

At discharge from therapy, patients are transitioned to a home-based or health club maintenance regimen. A recent study^[63] demonstrated high compliance with exercise recommendations at follow-up. For those with work disability, recommendations for return to work are based upon performance in therapy.

4. Studies Utilising Aggressive Exercise Regimens

Data from prospective and retrospective studies utilising aggressive exercise as treatment for patients with chronic low back pain reveal that within a period of 6 to 8 weeks, it is possible to improve trunk flexibility by 20%, trunk strength and lifting capacities by 50%, and endurance by 20 to 60%.^[15,16,20,37,46,64,65] Pain-related disability as determined by the Oswestry scale^[66] was reduced by 50%, on average, and pain severity by 30%.^[67] Treatment outcomes, once quantified, are amenable to analysis using statistical databases. By tracking treatment outcomes, information can be derived to modify treatment regimens, with the goal of improving patient care.

5. Conclusion

Chronic back pain is often a complex condition, influenced by pathology, pain intensity, impairments in back function, and societal and behavioural reinforcements, with significant resulting disability. To date, there is no scientific evidence that activity and exercises are harmful, or that pain-inducing activity must be avoided by this patient population. Indeed, empirical evidence to the contrary suggests that activity and exercise that challenge physical impairments actually result in an improvement in chronic back pain. The exercise philosophy and regimen outlined in this article exemplifies one such approach that combines aggressive exercise with focused application of cognitive behavioural techniques.

Clearly, future research is needed in this area. We need a better understanding of how exercise alters psychological fears and beliefs regarding pain and function. Dose-response relationships regarding the intensity of exercise and outcomes require further clarification. Predictors of outcome should be further explored, with the hope that this will lead to modification of therapy and improved outcomes for those most resistant to our care.

Acknowledgements

The authors have no conflicts of interest.

Appendix

Exercises for the treatment of low back pain.

Table A1. A summary of the exercises/machines and the specific muscle groups they target in the treatment of patients with low back pain

Exercise	Muscles
Cybox back extension	Multifidi, longissimus, iliocostalis
Roman chair hyperextension	Multifidi, longissimus, iliocostalis, gluteus maximus, hamstrings
Lumbar crate lifting	Multifidi, longissimus, iliocostalis, gluteus maximus, hamstrings, rhomboids, trapezius
Lat pull-down	Latissimus dorsi, rhomboids, teres major/minor
Cybox rotary torso	Internal/external obliques, adductors, erector spinae
Multihip machine	Hip flexors, hip extensors, hip abductors, hip adductors, glutei, erector spinae




Fig. A1. Examples of some exercises for the treatment of low back pain: (a) Cybex back extension; (b) lumbar crate lifting; (c) lat pulldown; and (d) Cybex multihip.

References

- Bigos SJ, Battie MC, Spengler DM, et al. A prospective study of work perceptions and psychosocial factors affecting the report of back injury. *Spine* 1991; 16: 1-6
- Frymoyer JW, Rosen JC, Clements J, et al. Psychologic factors in low-back-pain disability. *Clin Orthop* 1985 May; 195: 178-84
- Rainville J, Sobel JB, Hartigan C, et al. The effect of compensation involvement on the reporting of pain and disability by patients referred for rehabilitation of chronic low back pain. *Spine* 1997; 22: 2016-24
- Tollison CD. Compensation status as a predictor of outcome in non-surgically treated low back injury. *South Med J* 1993; 86: 1206-9
- Rainville J, Ahern DK, Phalen L. Altering beliefs about pain and impairment in a functionally oriented treatment program for chronic low back pain. *Clin J Pain* 1993; 9: 196-201
- Riley JF, Ahern DK, Follick MJ. Chronic pain and functional impairment: assessing beliefs about their relationship. *Arch Phys Med Rehabil* 1988; 69: 579-82
- Kori SH, Miller RP, Todd DD. Kinisophobia: a new view of pain behavior. *Pain Manage* 1990 Jan-Feb; 35-43

8. Waddell G, Newton M, Henderson I. A fear avoidance belief questionnaire (FABQ) and the role of fear avoidance beliefs in chronic low back pain. *Pain* 1993; 52: 158-6
9. Rainville J, Carlson N, Polatin P, et al. Exploration of physicians' recommendations for activities in chronic low back pain. *Spine* 2000; 25: 2210-20
10. Dolce JJ, Crocker MF, Molettire C, et al. Exercise quotas, anticipatory concerns and self efficacy in chronic pain: a preliminary report. *Pain* 1986; 24: 365-72
11. Rainville J, Bagnall D, Phalen L. Health care providers' attitudes and beliefs about functional impairments and chronic back pain. *Clin J Pain* 1995; 11: 287-95
12. Alaranta H, Rytokoski U, Rissanen A, et al. Intensive physical and psychosocial training program for patients: a controlled clinical trial. *Spine* 1994; 12: 1339-49
13. Bendix AF, Bendix T, Labriola M, et al. Functional restoration for chronic low back pain: two-year followup of two randomized clinical trials. *Spine* 1998; 23: 717-25
14. Carpenter DM, Nelson BW. Low back strengthening for the prevention and treatment of low back pain. *Med Sci Sports Exerc* 1999; 31: 18-24
15. Hazard RG, Fenwick JW, Kalisch SM, et al. Functional restoration with behavioral support: a one year prospective study of patients with chronic low back pain. *Spine* 1989; 14: 157-61
16. Mayer TG, Gatchel RJ, Kishino N, et al. 1985 Volvo Award in clinical sciences: Objective assessment of spine function following industrial injury - a prospective study with comparison group and one year follow-up. *Spine* 1985; 10: 482-93
17. Mayer TG, Gatchel RJ, Mayer H, et al. A prospective two year study of functional restoration in industrial low back injury: an objective assessment procedure. *JAMA* 1987; 258: 1763-7
18. Mitchell RI, Carmen GM. Results of a multicenter trial using an intensive active exercise program for the treatment of acute soft tissue and back injuries. *Spine* 1990; 15: 514-21
19. Oland G, Tveiten G. A trial of modern rehabilitation for chronic low back pain and disability: vocational outcome and effect of pain modulation. *Spine* 1991; 16: 457-9
20. Risch SV, Norvell NK, Pollock ML, et al. Lumbar strengthening in chronic low back pain patients: physiologic and psychological benefits. *Spine* 1993; 18: 232-8
21. Sachs BL, David JF, Olimpio D, et al. Spinal rehabilitation by work tolerance based on objective physical capacity assessment of dysfunction: a prospective study with control subjects and twelve month review. *Spine* 1990; 15: 1325-32
22. Chiarello CM, Savidge R. Interrater reliability of the Cybex EDI-320 and fluid goniometer in normals and patients with low back pain. *Arch Phys Med Rehabil* 1993; 74: 32-7
23. Keeley J, Mayer TG, Cox R, et al. Quantification of lumbar function. Part 5: reliability of range-of-motion measures in the sagittal plane and an *in vivo* torsion rotation measurement technique. *Spine* 1986; 11: 31-5
24. Mayer TG, Tencer AF, Kristopherson S, et al. Use of non-invasive techniques for quantification of spinal range of motion in normal subjects and chronic low back dysfunction patients. *Spine* 1984; 9: 588-95
25. Mayer TG, Tabor J, Bovasso E, et al. Physical progress and impairment quantification after functional restoration. Part I: lumbar mobility. *Spine* 1994; 19: 389-94
26. Waddell G, Somerville D, Henderson I, et al. Objective clinical evaluation of physical impairment in chronic low back pain. *Spine* 1992; 17: 617-28
27. Anderson JA, Sweetman BJ. A combined flexirule/hydrogoniometer for measurement of lumbar spine and its sagittal movement. *Rheum Rehabil* 1975; 14: 173-9
28. Burdett RG, Brown KE, Fall MP. Reliability and validity of four instruments for measuring lumbar spine and pelvic positions. *Phys Ther* 1986; 66: 677-84
29. Loebel W. Measurement of spinal posture and range of spinal movement. *Ann Phys Med* 1967; 9: 103-10
30. Macrae IF, Wright V. Measurement of back movement. *Ann Rheum Dis* 1969; 28: 584-9
31. Mooney V. Physical measurements of the lumbar spine. *Phys Med Rehabil Clin N Am* 1998; 9: 391-408
32. Tanz SS. Motion of the lumbar spine: a roentgenologic study. *Am J Roentgenol Rad Ther Nucl Med* 1953; 69: 399-412
33. Troup LT, Hood C, Chapman AE. Measurements of sagittal mobility of the lumbar spine and hips. *Ann Phys Med* 1968; 9: 308-21
34. Rainville J, Sobel JB, Hartigan C. Comparison of total lumbosacral flexion and true lumbar flexion measured by a dual inclinometer technique. *Spine* 1994; 19: 2698-701
35. Fitzgerald GK, Wynveen KJ, Rheault W, et al. Objective assessment with establishment of normal values for lumbar spinal motion. *Phys Ther* 1983; 63: 1776-81
36. Addison R, Schultz A. Trunk strengths in patients seeking hospitalization for chronic low back pain disorders. *Spine* 1980; 5: 539-44
37. Brady S, Mayer TG, Gatchel RJ. Physical progress and residual impairment quantification after functional restoration. Part II: isokinetic trunk strength. *Spine* 1994; 19: 395-400
38. Kahanovitz N, Viola K, Gallagher M. Long term strength assessment of postoperative discectomy patients. *Spine* 1989; 14: 402-3
39. Mayer TG, Smith SS, Keeley J, et al. Quantification of lumbar function. Part 2: sagittal plane trunk strength in chronic-low back pain patients. *Spine* 1985; 10: 765-72
40. Mayer TG, Vanharanta H, Gatchel RJ, et al. Comparison of CT scan muscle measurements and iso-kinetic trunk strength in postoperative patients. *Spine* 1989; 14: 33-6
41. McNeill T, Warwick D, Andersson GBJ, et al. Trunk strengths in attempted flexion, extension and lateral bending in healthy subjects and patients with low back pain. *Spine* 1980; 5 (6): 529-38
42. Nachemson AL, Lindh M. Measurement of abdominal and back muscle strength with and without low back pain. *Scand J Rehabil Med* 1969; 1: 60-5
43. Novy DM, Simmonds MJ, Olson SL, et al. Physical performance: differences in men and women with and without low back pain. *Arch Phys Med Rehabil* 1999; 80: 195-8
44. Smith SS, Mayer TG, Gatchel RJ, et al. Quantification of lumbar function. Part I: isometric and multispeed isokinetic trunk strength measures in sagittal and axial planes in normal subjects. *Spine* 1985; 10: 757-64
45. Rainville J, Sobel J, Hartigan C, et al. Decreasing disability in chronic back pain through aggressive spine rehabilitation. *J Rehabil Res Dev* 1997; 34: 383-93
46. Curtis L, Mayer TG, Gatchel RJ. Physical progress and residual impairment quantification after functional restoration. Part III: isokinetic and isoinertial lifting capacity. *Spine* 1994; 19: 401-5
47. Kishino ND, Mayer TG, Gatchel RJ, et al. Quantification of lumbar function. Part 4: isometric and isokinetic lifting simulation in normal subjects and low back dysfunction patients. *Spine* 1985; 10: 921-7

48. Mayer TG, Barnes D, Kishino ND, et al. Progressive isoinertional lifting evaluation. Part 1: a standard protocol and normative database. *Spine* 1988; 13: 993-7
49. Bortz WM. The disuse syndrome. *West J Med* 1984; 141: 691-4
50. Kashihara H, Haruna Y, Suzuki Y, et al. Effects of mild supine exercise during 20 days bedrest on maximal oxygen uptake rate in young humans. *Acta Physiol Scand Suppl* 1994; 616: 19-26
51. Kottke FJ. The effects of limitation of activity upon the human body. *JAMA* 1966; 196: 825-30
52. Saltin B, Blomquist G, Mitchell J, et al. Response to exercise after bedrest and after training: a longitudinal study of adaptive changes in oxygen transport and composition. *Circulation* 1968; 38 Suppl. 7: 1-78
53. Takenaka F, Suzuki Y, Kawakubo K, et al. Cardiovascular effects of 20 days bedrest in healthy young subjects. *Acta Physiol Scand Suppl* 1994; 616: 59-63
54. Boumfrey FR, Levenberg RJ. A prospective study of the role of anaerobic conditioning in the treatment of chronic low back pain. 9th Annual Meeting of the North American Spine Society; 1994 Oct 19-20; Minneapolis
55. Browman CP. Sleep following sustained exercise. *Psychophysiology* 1980; 17: 577-80
56. DeVries H, Adams GM. EMG comparisons of single doses of exercise and meprobamate as to effects on muscle relaxation. *Am J Phys Med* 1972; 51: 130-41
57. DeVries H, Wiswell RA, Bulbion R, et al. Tranquilizer effects of exercise: acute effects of moderate aerobic exercise on spinal reflex activation level. *Am J Phys Med* 1981; 60: 57-66
58. Mellion MB. Exercise therapy for anxiety and depression. Part 1: does the evidence justify its recommendation? *Postgrad Med* 1985; 77: 59-62, 66
59. Tollison CD, Kriegel ML, Downie GR. Chronic low back pain: results of treatment at the Pain Therapy Center. *South Med J* 1985; 78: 1291-5
60. Schmidt AJ. Cognitive factors in the performance level of chronic low back pain patients. *J Psychosom Res* 1985; 29: 183-9
61. Fordyce WE, Shelton JL, Dunmore DE. The modification of avoidance learning pain behaviors. *J Behav Med* 1982; 5: 405-14
62. Waddell G, Main CJ, Morris EW, et al. Chronic low back pain, psychologic distress, and illness behavior. *Spine* 1984; 9: 209-13
63. Hartigan C, Rainville J, Sobel JB, et al. Long-term exercise adherence after intensive rehabilitation for chronic low back pain. *Med Sci Sports Exerc* 2000; 32: 551-7
64. Mayer T, Tabor J, Bovasso E, et al. Physical progress and residual impairment quantification after functional restoration. Part I: lumbar mobility. *Spine* 1994; 19: 389-94
65. Pollock ML, Leggett SH, Graves JE, et al. Effect of resistance training on lumbar extension strength. *Am J Sports Med* 1989; 17: 624-9
66. Fairbank JCT, Couper J, Davis JB, et al. The Oswestry low back pain disability questionnaire. *Physiotherapy* 1980; 66: 271-3
67. Hartigan C, Sobel JB, Rainville J. Functionally oriented rehabilitation in low back pain: changes in pain scores. 9th Annual Meeting of the North American Spine Society; 1994 Oct 19-20; Minneapolis

Correspondence and offprints: Dr *James Rainville*, The Spine Center, New England Baptist Hospital, 125 Parker Hill Avenue, Boston, MA 02120, USA.
E-mail: jrainvil@caregroup.harvard.edu